

When new transgenic lines need to be introduced into animal facilities, or when animals need to be rederived to remove specific pathogens, this Shared Resource is capable of performing a variety of techniques necessary to prevent the introduction of pathogens in the mouse facilities. This includes *in vitro* fertilization, rederivation and ovarian transplantation. Specific Pathogen Free animals will be generated and delivered to your facility.

Keywords: [rederivation](#) [TMESCSR](#) [service](#) [reproduction](#) [mouse](#) [mice](#) [IVF](#) [embryo transfer](#) [in vitro fertilization](#)

[Expand](#)

***In vitro* fertilization (IVF)**

In vitro fertilization requires fresh or frozen sperm from a transgenic male that is used to fertilize oocytes from superovulated female mice. IVF rederivations generally produce larger numbers of pups born when compared to the thawing of one or two straws of embryos for rederivation. After incubation overnight and fertilization, two-cell embryos are transferred into recipient animals. About 19 days later, pups are born and at 3 weeks of age the Vanderbilt Genome Editing Resource staff will wean, tail, and ear punch the pups. The investigator will then screen the tail DNA for transgenic founders and notify the resource of their results. This is also a valuable service for quick expansion of a mouse colony, providing a large quantity of animals to establish a colony.

Embryo Transfer Rederivation

For rederivations, fresh or frozen 0.5 to 3.5 day old embryos are washed numerous times in sterile medium and are then transferred into recipient females. Seventeen to nineteen days later pups are born and at 3 weeks of age the Vanderbilt Genome Editing Resource staff will wean, tail, and ear punch the pups. The investigator will then screen the tail DNA for transgenic founders and notify the resource of their results. This service is recommended for homozygous lines or lines with multiple genetic mutations.

Embryo Retrieval and Transfer Rederivation

For embryo transfer, females are superovulated and bred to males. Embryos are collected the following day, washed to further reduce pathogen transfer and then surgically transferred into pseudopregnant females housed in the barrier facility. Pups will be born about 19 days later and weaned and tailed at 3 weeks of age. This service is recommended for homozygous lines or lines with multiple genetic mutations.

Ovary Transplants

For ovary transplantation, ovaries from a transgenic female are removed and transplanted into 1-2 recipient animals. This service, although not frequently utilized, is essential when a valuable line of animals ceases to reproduce and is at risk.

***In Vitro* fertilization**


Rederivation

Ovary transplantation


- | | | |
|---|--|--|
| <ol style="list-style-type: none">1. PI submits appropriate service form to VGER2. Service forms are reviewed and approved by the Co-Directors of the resource3. IVF is scheduled, mice are ordered4. IVF is performed5. The following day, fertilized embryos are transferred to pseudopregnant females6. Pups born 19 to 20 days post fertilization7. Pups weaned, tailed and ear punched by VGER staff at 3 weeks of age8. DAC staff review serology results, approve movement of mice, and transfer transgenic founders to PIs mouse housing room. | <ol style="list-style-type: none">1. PI submits appropriate service form to VGER2. Service forms are reviewed and approved by the Co-Directors of the resource3. Rederivation is scheduled, mice are ordered (if required)4. Rederivation is performed, embryos are transferred to pseudopregnant females5. Pups born 17 to 20 days following transfer6. Pups weaned, tailed and ear punched by VGER staff at 3 weeks of age7. PI identifies transgenic founders from tail DNA using PCR and Southern blotting8. PI notifies VGER staff about transgenic founders9. DAC staff review serology results, approve movement of mice, and transfer transgenic founders to PIs mouse housing room. | <ol style="list-style-type: none">1. PI submits appropriate service form to VGER2. Service forms are reviewed and approved by the Co-Directors of the resource3. Ovary transplantation is scheduled4. Ovary transplant is performed5. Mice will be housed in the barrier facility for 4 weeks6. Females will be serology tested after 4 weeks7. DAC staff review serology results, approve movement of mice, and perform transfer of mice to PI's mouse housing room |
|---|--|--|


[Click here to contact the core](#)

Attachments

 [IVF_Dishes.jpg](#) - Added on March 18, 2016 at 2:48 PM by [Jennifer Skelton](#)

IVF dishes sitting in a modular incubator.



 [Rederivation_Service_Form_0119.docx](#) - Added on December 18, 2018 at 12:03 PM by [Jennifer Skelton](#)